

## Post-Operative Instructions: Lymph Node Biopsy

H. Dennis Snyder, M.D., F.A.C.S.

Seema V. Byahatti, M.D., F.A.C.S., MBA

Russell S. Shu, M.D., F.A.C.S.

Douglas A. O'Brien, M.D.

Pratik S. Pradhan, M.D.

Jagdish K. Dhingra, M.D., F.R.C.S.

Tiffany A. Hron, M.D.

Walid I. Dagher, M.D.

[entspecialists.com](http://entspecialists.com)

### **Norwood**

825 Washington Street, Suite 310

MA, 02062

P: 781-769-8910

F: 781-255-9844

### **Brockton**

35 Pearl Street, Suite 200

MA, 02301

P: 508-588-8034

F: 508-897-0475

### **Taunton**

72 Washington Street, Suite 1600

MA, 02780

P: 508-880-3460

F: 508-880-5335

### **Plainville**

188 Washington Street

MA, 02762

P: 508-699-1701

F: 508-699-1706

- **Diet:** Drink plenty of fluids to keep yourself well hydrated. You may eat your regular diet.
- **Pain:** Pain medication will be prescribed by your doctor, take as directed. You may also take Tylenol® or Ibuprofen for pain. Do not take any aspirin-containing medications for 2 weeks after surgery. Check medication labels carefully.
- **Swelling:** It is possible for fluid and/or blood to accumulate under the skin at the surgical site. If you notice bulging and excessive swelling in this area, please contact your doctor.
- **Fever:** You may experience a low-grade fever (100-101°F) for 2-3 days after surgery. If a fever persists beyond this time, contact your doctor.
- **Activity:** No strenuous activity, heavy lifting, swimming, or exercise for 2 weeks after surgery. You may return to work in 7-10 days, but no strenuous activity or lifting at work for 2 weeks.
- **Wound Care:** Apply Bacitracin or Vaseline® ointment over your sutures twice daily, unless otherwise directed by your doctor. Wash your hands before applying ointment. Keep the incision line dry for 48 hours after surgery, after which you may shower. After showering, carefully dry the incision line and apply ointment.

For patients who take Coumadin® or any other blood thinner, check with your primary care doctor for instructions as to when to resume these medications after surgery.